


GEOLOGIC MAP OF THE THOUSAND SPRINGS QUADRANGLE, GOODING AND TWIN FALLS COUNTIES, IDAHO

Virginia S. Gillerman, John D. Kauffman, and Kurt L. Othberg

Disclaimer: This Digital Web Map is an informal report and may be revised and formally published at a later time. Its content and format may not conform to agency standards.

2005



MAP SYMBOLS

- Contact: Line showing the approximate boundary between one map unit and another. The location accuracy of an approximate contact is more than 80 feet on the ground.
- Fault: Approximately located; dotted where concealed; queried where uncertain; ball and bar on downthrown side.
- Baked sediments in contact with basalt flows.
- Bonneville Flood flow direction.
- Location of evidence for maximum stage of Bonneville Flood. Numbered location reported by O'Connor (1993).
- Approximate extent of Bonneville Flood at maximum stage.
- Trend of dune field: arrow points in the downwind direction.
- Stabilized dune field.
- Landslide scarp and headwall: ticks show top of scarp.
- Gravel pit that exposes a map unit.
- Monoclinical fold: Dashed were approximately located; dotted where concealed; queried where uncertain.
- Strike and dip of basalt flow or sedimentary bedding.
- Estimated strike and dip of basalt flow.
- Sample site for chemical and paleomagnetic analyses.*
- Sample site for chemical analysis.*
- Sample site for paleomagnetic analysis.*

INTRODUCTION


The geologic map of the Thousand Springs quadrangle identifies both the bedrock and surficial geologic units. It shows the geographic distribution of rock types at the surface and in the shallow subsurface. The geologic units in the area control soil development, groundwater flow, and the distribution of agricultural enterprises, and dairy farms with confined animal feeding operations. In addition, a structurally controlled geothermal system exists in the western portion of the quadrangle and provides commercial hot water for geothermal enterprises, and dairy farms with confined animal feeding operations. In addition, a structurally controlled geothermal system exists in the western portion of the quadrangle and provides commercial hot water for geothermal enterprises, and dairy farms with confined animal feeding operations. In addition, a structurally controlled geothermal system exists in the western portion of the quadrangle and provides commercial hot water for geothermal enterprises, and dairy farms with confined animal feeding operations.

DESCRIPTION OF MAP UNITS

ARTIFICIAL DEPOSITS

Made ground (Holocene)—Artificial fills composed of excavated, transported, and/or compacted materials typically derived locally. Primarily areas modified for fish ponds.

CORRELATION OF MAP UNITS



ALLUVIAL AND LACUSTRINE DEPOSITS

Qam Alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel of channel bars, islands, and shorelines. Gravelly where channel is shallow and formed directly in basalt. In Salmon Falls Creek canyon includes terrace and alluvial fan deposits. Typically 1-10 feet thick.

Qoa Older alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel in Melon Valley.

Qas Alluvium of side-streams (Holocene)—Channel and flood-plain deposits of tributaries to the Snake River. Stratified silt, sand, and gravel in Melon Valley.

Qab Alluvial fan deposits (Holocene)—Stratified silt, sand, and gravel that form small fans adjacent to Salmon Falls Creek. Merges and is interstratified with alluvium (Qam, Qas). Thickness varies, but typically ranges 3-30 feet.

Qag Sand and gravel in giant flood bars (Pleistocene)—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlines with rapid expansion basins with large-scale crossbeds. Deposited during high-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qaf Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)—Stratified coarse sand and pebble-cobble gravel deposited in eddy, side-channel positions, and lower-energy, waving-stage flood channels. Mantled with thin lens and minor fine-grained alluvium and slope wash.

Qah Scabland of flood pathways (Pleistocene)—Flood-scoured basalt surface. Above the canyon rim, scoured surface is stripped of pre-flood soils but thin post-flood lenses and soil are discontinuously present. In the canyon, sedimentary cover has been stripped and basalt surfaces have been panned, gouged, and smoothed. Includes minor deposits of coarse sand that are not mapped at this scale. Some areas include pavements or strings of boulders transported by flood traction forces or that are lags from erosion by lower-energy regime during late stages of the flood.

Qai Crowned Gravel (Pleistocene)—Stratified sand and pebble gravel that overlies Yabou Clay (Qy). Gravel clasts composed of felsic volcanic rocks, quartzite, chert, and minor basalt. Map location suggests unit is channel deposits of ancestral Snake River that prograded across Yabou Clay as McKinney Lake regressed (see Qy). Thickness about 6 feet. Original thickness and extent unknown owing to erosion by Bonneville Flood.

Qaj Sand and gravel in giant flood bars (Pleistocene)—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlines with rapid expansion basins with large-scale crossbeds. Deposited during high-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qak Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)—Stratified coarse sand and pebble-cobble gravel deposited in eddy, side-channel positions, and lower-energy, waving-stage flood channels. Mantled with thin lens and minor fine-grained alluvium and slope wash.

Qal Scabland of flood pathways (Pleistocene)—Flood-scoured basalt surface. Above the canyon rim, scoured surface is stripped of pre-flood soils but thin post-flood lenses and soil are discontinuously present. In the canyon, sedimentary cover has been stripped and basalt surfaces have been panned, gouged, and smoothed. Includes minor deposits of coarse sand that are not mapped at this scale. Some areas include pavements or strings of boulders transported by flood traction forces or that are lags from erosion by lower-energy regime during late stages of the flood.

Qam Alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel of channel bars, islands, and shorelines. Gravelly where channel is shallow and formed directly in basalt. In Salmon Falls Creek canyon includes terrace and alluvial fan deposits. Typically 1-10 feet thick.

Qoa Older alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel in Melon Valley.

Qas Alluvium of side-streams (Holocene)—Channel and flood-plain deposits of tributaries to the Snake River. Stratified silt, sand, and gravel in Melon Valley.

Qab Alluvial fan deposits (Holocene)—Stratified silt, sand, and gravel that form small fans adjacent to Salmon Falls Creek. Merges and is interstratified with alluvium (Qam, Qas). Thickness varies, but typically ranges 3-30 feet.

Qag Sand and gravel in giant flood bars (Pleistocene)—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlines with rapid expansion basins with large-scale crossbeds. Deposited during high-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qaf Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)—Stratified coarse sand and pebble-cobble gravel deposited in eddy, side-channel positions, and lower-energy, waving-stage flood channels. Mantled with thin lens and minor fine-grained alluvium and slope wash.

Qah Scabland of flood pathways (Pleistocene)—Flood-scoured basalt surface. Above the canyon rim, scoured surface is stripped of pre-flood soils but thin post-flood lenses and soil are discontinuously present. In the canyon, sedimentary cover has been stripped and basalt surfaces have been panned, gouged, and smoothed. Includes minor deposits of coarse sand that are not mapped at this scale. Some areas include pavements or strings of boulders transported by flood traction forces or that are lags from erosion by lower-energy regime during late stages of the flood.

Qai Crowned Gravel (Pleistocene)—Stratified sand and pebble gravel that overlies Yabou Clay (Qy). Gravel clasts composed of felsic volcanic rocks, quartzite, chert, and minor basalt. Map location suggests unit is channel deposits of ancestral Snake River that prograded across Yabou Clay as McKinney Lake regressed (see Qy). Thickness about 6 feet. Original thickness and extent unknown owing to erosion by Bonneville Flood.

Qaj Sand and gravel in giant flood bars (Pleistocene)—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlines with rapid expansion basins with large-scale crossbeds. Deposited during high-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qak Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)—Stratified coarse sand and pebble-cobble gravel deposited in eddy, side-channel positions, and lower-energy, waving-stage flood channels. Mantled with thin lens and minor fine-grained alluvium and slope wash.

Qal Scabland of flood pathways (Pleistocene)—Flood-scoured basalt surface. Above the canyon rim, scoured surface is stripped of pre-flood soils but thin post-flood lenses and soil are discontinuously present. In the canyon, sedimentary cover has been stripped and basalt surfaces have been panned, gouged, and smoothed. Includes minor deposits of coarse sand that are not mapped at this scale. Some areas include pavements or strings of boulders transported by flood traction forces or that are lags from erosion by lower-energy regime during late stages of the flood.

Qam Alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel of channel bars, islands, and shorelines. Gravelly where channel is shallow and formed directly in basalt. In Salmon Falls Creek canyon includes terrace and alluvial fan deposits. Typically 1-10 feet thick.

Qoa Older alluvium of mainstreams (Holocene)—Channel and flood-plain deposits of the Snake River and Salmon Falls Creek. Stratified silt, sand, and gravel in Melon Valley.

Qas Alluvium of side-streams (Holocene)—Channel and flood-plain deposits of tributaries to the Snake River. Stratified silt, sand, and gravel in Melon Valley.

Qab Alluvial fan deposits (Holocene)—Stratified silt, sand, and gravel that form small fans adjacent to Salmon Falls Creek. Merges and is interstratified with alluvium (Qam, Qas). Thickness varies, but typically ranges 3-30 feet.

Qag Sand and gravel in giant flood bars (Pleistocene)—Stratified deposits of boulders, cobbles, and pebbles of basalt in a matrix of coarse sand. Forms streamlines with rapid expansion basins with large-scale crossbeds. Deposited during high-energy, maximum stage of flood. Similar to Melon Gravel (Malde and Powers, 1962; Malde and others, 1963; and Covington and Weaver, 1991), but restricted to Bonneville Flood constructional forms and deposits.

Qaf Sand and gravel in eddy deposits and lower-energy bars (Pleistocene)—Stratified coarse sand and pebble-cobble gravel deposited in eddy, side-channel positions, and lower-energy, waving-stage flood channels. Mantled with thin lens and minor fine-grained alluvium and slope wash.

Qah Scabland of flood pathways (Pleistocene)—Flood-scoured basalt surface. Above the canyon rim, scoured surface is stripped of pre-flood soils but thin post-flood lenses and soil are discontinuously present. In the canyon, sedimentary cover has been stripped and basalt surfaces have been panned, gouged, and smoothed. Includes minor deposits of coarse sand that are not mapped at this scale. Some areas include pavements or strings of boulders transported by flood traction forces or that are lags from erosion by lower-energy regime during late stages of the flood.

Qai Crowned Gravel (Pleistocene)—Stratified sand and pebble gravel that overlies Yabou Clay (Qy). Gravel clasts composed of felsic volcanic rocks, quartzite, chert, and minor basalt. Map location suggests unit is channel deposits of ancestral Snake River that prograded across Yabou Clay as McKinney Lake regressed (see Qy). Thickness about 6 feet. Original thickness and extent unknown owing to erosion by Bonneville Flood.

BASALT UNITS

The surficial geology of the Snake River Plain north of the Snake River is primarily Pleistocene basalt flows of the Snake River Group. North of the Snake River, the younger basalt units originated from several shield volcanoes north and east of the quadrangle, whereas older Pliocene and Miocene basalt units originated south of the Snake River. Each volcano probably extruded numerous lava flows or flow lobes, although individual flows cannot easily be mapped, especially on the older surfaces now subdued by surficial deposits. Older basalt surfaces tend to be less rugged and more subdued than younger surfaces, primarily the result of greater accumulation of loss over a longer period of time.

Qib Basalt of North Butte (Pleistocene)—Fine-grained, dark gray to black basalt with common to abundant olivine phenocrysts and clots 0.5 to 1.5 mm in diameter. Locally contains a few small plagioclase phenocrysts as much as 1 mm in length, or scarce to rare glomerulites of plagioclase and olivine as much as 5 mm. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is North Butte located 2.1 miles northeast of the quadrangle. A small lobe extends onto the quadrangle from the north. Equivalent to the Wendell Grade Basalt of Malde and Powers (1962) and Malde and others (1963).

Qic Basalt of Bacon Butte, younger unit (Pleistocene)—Fine- to medium-grained, dark gray basalt with common to abundant olivine as grains and clots as much as 3 mm and scattered small plagioclase laths. Similar in texture and appearance to basalt of North Butte, although slightly coarser grained overall. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Forms a raised, hilly surface of partly exposed pressure ridges that trends from east-to-west into the Sand Springs area in the Snake River canyon, where Malde and others (1963) mapped as Sand Springs Basalt. The surface was scoured where the Bonneville Flood crossed the upland, rendering an appearance that could be mistaken for a younger basalt, such as basalt of Rocky Butte. However, we believe this unit is a late flow or series of flows erupted from Bacon Butte located 18 miles northeast of the quadrangle, which was named for "Bacon Ranch." The name "Bacon Butte" is derived from nearby Bacon Ranch, located on the east side of the butte. The name was also used by Covington and Weaver (1991). Surface drainage is not developed to poorly developed. Discontinuous lens and columnar sand deposits occur less than 50 percent of the surface and are 1-10 feet thick. Soil caliche (duripan) is commonly well developed within the soil profile (Young and others, 1929; Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche varies considerably. Some of the land is cultivatable and some is used as pasture.

Qid Basalt of Bacon Butte, older unit (Pleistocene)—Fine-grained, dark gray basalt with common to abundant plagioclase laths as much as 5 mm in length and common olivine grains and clots, commonly as intergrowths with plagioclase. Locally diatrytic. May exhibit abundant carbonate accumulation in vesicles and fractures. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Also erupted from Bacon Butte located 18 miles northeast of the quadrangle. Included in Thousand Springs Basalt by Malde and others (1963). Topography contains with areas of basalt of North Butte and the basalt of Bacon Butte, younger unit. Few basalt pressure ridges rise above a nearly complete mantle of loss and dune sand. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche is highly variable. Most of the land is cultivatable.

Qie Basalt of Flat Top Butte (Pleistocene)—Fine-grained, medium gray basalt with scattered to very abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Flat Top Butte located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qif Basalt of Burger Butte (late Pliocene)—Fine- to medium-grained basalt generally with abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Burger Butte and associated satellite vents located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qig Basalt of North Butte (Pleistocene)—Fine-grained, dark gray to black basalt with common to abundant olivine phenocrysts and clots 0.5 to 1.5 mm in diameter. Locally contains a few small plagioclase phenocrysts as much as 1 mm in length, or scarce to rare glomerulites of plagioclase and olivine as much as 5 mm. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is North Butte located 2.1 miles northeast of the quadrangle. A small lobe extends onto the quadrangle from the north. Equivalent to the Wendell Grade Basalt of Malde and Powers (1962) and Malde and others (1963).

Qih Basalt of Bacon Butte, younger unit (Pleistocene)—Fine- to medium-grained, dark gray basalt with common to abundant olivine as grains and clots as much as 3 mm and scattered small plagioclase laths. Similar in texture and appearance to basalt of North Butte, although slightly coarser grained overall. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Forms a raised, hilly surface of partly exposed pressure ridges that trends from east-to-west into the Sand Springs area in the Snake River canyon, where Malde and others (1963) mapped as Sand Springs Basalt. The surface was scoured where the Bonneville Flood crossed the upland, rendering an appearance that could be mistaken for a younger basalt, such as basalt of Rocky Butte. However, we believe this unit is a late flow or series of flows erupted from Bacon Butte located 18 miles northeast of the quadrangle, which was named for "Bacon Ranch." The name "Bacon Butte" is derived from nearby Bacon Ranch, located on the east side of the butte. The name was also used by Covington and Weaver (1991). Surface drainage is not developed to poorly developed. Discontinuous lens and columnar sand deposits occur less than 50 percent of the surface and are 1-10 feet thick. Soil caliche (duripan) is commonly well developed within the soil profile (Young and others, 1929; Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche varies considerably. Some of the land is cultivatable and some is used as pasture.

Qii Basalt of Bacon Butte, older unit (Pleistocene)—Fine-grained, dark gray basalt with common to abundant plagioclase laths as much as 5 mm in length and common olivine grains and clots, commonly as intergrowths with plagioclase. Locally diatrytic. May exhibit abundant carbonate accumulation in vesicles and fractures. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Also erupted from Bacon Butte located 18 miles northeast of the quadrangle. Included in Thousand Springs Basalt by Malde and others (1963). Topography contains with areas of basalt of North Butte and the basalt of Bacon Butte, younger unit. Few basalt pressure ridges rise above a nearly complete mantle of loss and dune sand. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche is highly variable. Most of the land is cultivatable.

Qij Basalt of Flat Top Butte (Pleistocene)—Fine-grained, medium gray basalt with scattered to very abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Flat Top Butte located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qik Basalt of Burger Butte (late Pliocene)—Fine- to medium-grained basalt generally with abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Burger Butte and associated satellite vents located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qil Basalt of North Butte (Pleistocene)—Fine-grained, dark gray to black basalt with common to abundant olivine phenocrysts and clots 0.5 to 1.5 mm in diameter. Locally contains a few small plagioclase phenocrysts as much as 1 mm in length, or scarce to rare glomerulites of plagioclase and olivine as much as 5 mm. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is North Butte located 2.1 miles northeast of the quadrangle. A small lobe extends onto the quadrangle from the north. Equivalent to the Wendell Grade Basalt of Malde and Powers (1962) and Malde and others (1963).

Qim Basalt of Bacon Butte, younger unit (Pleistocene)—Fine- to medium-grained, dark gray basalt with common to abundant olivine as grains and clots as much as 3 mm and scattered small plagioclase laths. Similar in texture and appearance to basalt of North Butte, although slightly coarser grained overall. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Forms a raised, hilly surface of partly exposed pressure ridges that trends from east-to-west into the Sand Springs area in the Snake River canyon, where Malde and others (1963) mapped as Sand Springs Basalt. The surface was scoured where the Bonneville Flood crossed the upland, rendering an appearance that could be mistaken for a younger basalt, such as basalt of Rocky Butte. However, we believe this unit is a late flow or series of flows erupted from Bacon Butte located 18 miles northeast of the quadrangle, which was named for "Bacon Ranch." The name "Bacon Butte" is derived from nearby Bacon Ranch, located on the east side of the butte. The name was also used by Covington and Weaver (1991). Surface drainage is not developed to poorly developed. Discontinuous lens and columnar sand deposits occur less than 50 percent of the surface and are 1-10 feet thick. Soil caliche (duripan) is commonly well developed within the soil profile (Young and others, 1929; Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche varies considerably. Some of the land is cultivatable and some is used as pasture.

Qin Basalt of Bacon Butte, older unit (Pleistocene)—Fine-grained, dark gray basalt with common to abundant plagioclase laths as much as 5 mm in length and common olivine grains and clots, commonly as intergrowths with plagioclase. Locally diatrytic. May exhibit abundant carbonate accumulation in vesicles and fractures. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Also erupted from Bacon Butte located 18 miles northeast of the quadrangle. Included in Thousand Springs Basalt by Malde and others (1963). Topography contains with areas of basalt of North Butte and the basalt of Bacon Butte, younger unit. Few basalt pressure ridges rise above a nearly complete mantle of loss and dune sand. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche is highly variable. Most of the land is cultivatable.

Qio Basalt of Flat Top Butte (Pleistocene)—Fine-grained, medium gray basalt with scattered to very abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Flat Top Butte located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qip Basalt of Burger Butte (late Pliocene)—Fine- to medium-grained basalt generally with abundant plagioclase-olivine intergrowths as large as 4-7 mm and olivine phenocrysts about 1 mm in diameter. Remnant magnetic polarity is reverse as determined in the field and through laboratory analysis. Source is Burger Butte and associated satellite vents located 16 miles southeast of the quadrangle. Most of the unit is equivalent to the "basalt of Sucker Flat" unit of Bonnichsen and Godchaux (1997b) in the adjacent Bull quadrangle. No basal pressure ridges rise above less mantle. Surface drainage is moderately well developed. Loss thickness ranges 5-25 feet and typically comprises a younger deposit with weak soil development and an underlying older lens with a thick caliche (duripan) horizon (Baldwin, 1925; Ames, 2003).

Qiq Basalt of North Butte (Pleistocene)—Fine-grained, dark gray to black basalt with common to abundant olivine phenocrysts and clots 0.5 to 1.5 mm in diameter. Locally contains a few small plagioclase phenocrysts as much as 1 mm in length, or scarce to rare glomerulites of plagioclase and olivine as much as 5 mm. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Source is North Butte located 2.1 miles northeast of the quadrangle. A small lobe extends onto the quadrangle from the north. Equivalent to the Wendell Grade Basalt of Malde and Powers (1962) and Malde and others (1963).

Qir Basalt of Bacon Butte, younger unit (Pleistocene)—Fine- to medium-grained, dark gray basalt with common to abundant olivine as grains and clots as much as 3 mm and scattered small plagioclase laths. Similar in texture and appearance to basalt of North Butte, although slightly coarser grained overall. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Forms a raised, hilly surface of partly exposed pressure ridges that trends from east-to-west into the Sand Springs area in the Snake River canyon, where Malde and others (1963) mapped as Sand Springs Basalt. The surface was scoured where the Bonneville Flood crossed the upland, rendering an appearance that could be mistaken for a younger basalt, such as basalt of Rocky Butte. However, we believe this unit is a late flow or series of flows erupted from Bacon Butte located 18 miles northeast of the quadrangle, which was named for "Bacon Ranch." The name "Bacon Butte" is derived from nearby Bacon Ranch, located on the east side of the butte. The name was also used by Covington and Weaver (1991). Surface drainage is not developed to poorly developed. Discontinuous lens and columnar sand deposits occur less than 50 percent of the surface and are 1-10 feet thick. Soil caliche (duripan) is commonly well developed within the soil profile (Young and others, 1929; Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche varies considerably. Some of the land is cultivatable and some is used as pasture.

Qis Basalt of Bacon Butte, older unit (Pleistocene)—Fine-grained, dark gray basalt with common to abundant plagioclase laths as much as 5 mm in length and common olivine grains and clots, commonly as intergrowths with plagioclase. Locally diatrytic. May exhibit abundant carbonate accumulation in vesicles and fractures. Remnant magnetic polarity is normal, as determined in the field and through laboratory analysis. Also erupted from Bacon Butte located 18 miles northeast of the quadrangle. Included in Thousand Springs Basalt by Malde and others (1963). Topography contains with areas of basalt of North Butte and the basalt of Bacon Butte, younger unit. Few basalt pressure ridges rise above a nearly complete mantle of loss and dune sand. Surface drainage is moderately developed. Thickness of mantle ranges 3-25 feet commonly 3-12 feet thick. Soil caliche (duripan) is typically well developed within the soil profile (Johnson, 2002) and at the soil-basalt contact, but the thickness of caliche is highly variable. Most of the land is cultivatable.

Qit Basalt of Flat Top Butte (Pleistocene)—Fine-grained, medium gray basalt with scattered to very abundant plagioclase-olivine intergrowths as